arXiv:1211.6835v2 [hep-th] 4 Feb 2013

Shearing Black Holes and Scans of the Quark Matter Phase Diagram

Brett McInnes

NORDITA, KTH Royal Institute of Technology and Stockholm University

Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden

and

National University of Singapore[1]

email: matmcinn@nus.edu.sg

ABSTRACT

Basic kinematic considerations strongly suggest that effects arising from quark polarization should be seen in peripheral heavy ion collisions producing very high angular momentum densities. No such effects have yet been observed in collisions corresponding to regions of the quark matter phase diagram where the quark chemical potential is small, but there is reason to hope that they may be in future. Using the holography of planar AdS black holes with angular momentum and electric charge, we suggest that the time available for these effects to develop varies quite rapidly across the quark matter phase diagram, in a way which might, in principle, be investigated through heavy ion collisions at future facilities such as FAIR and NICA.

---

[1] Permanent Address

1. Quark Polarization and AdS Black Holes

The quark matter phase diagram [1][2], representing states of quark matter at various values of the temperature and quark chemical potential, is currently understood only in an approximate or qualitative sense. Until recently, experimental investigations, based on the study of heavy ion collisions, have only explored the region of the diagram near to the temperature axis; but a move towards larger values of the quark chemical potential is in its early stages in the beam energy scan programme of the RHIC collider [3][4], and will reach maturity with the completion of the FAIR [5] and NICA [6] facilities.



Figure 1: Conjectured Quark Matter Phase Diagram

In broad terms, one version[2] of the phase diagram has the following structure. As in more familiar examples of phase diagrams, there are several lines separating various phases, with the Quark-Gluon Plasma [QGP] occupying all regions at sufficiently high temperature, and hadronic matter being found at lower temperatures and low values of the chemical potential. There is thought to be a critical point at a temperature T_c given approximately by $T_c \approx 175$ MeV and at a value of the quark chemical potential μ which is unknown but estimated to be in the range 1 - 4 times T_c. The phase line separating the QGP and hadronic phases is thought to bend *downwards* from this point [8][9], so that the QGP can exist, at relatively high values of μ, at substantially *lower* temperatures than would be possible at or near the $\mu = 0$ axis. The phase line continues downward until it reaches a triple point, at a location which is unknown; the nature of the third phase [which may in fact consist of several distinct phases] is likewise unknown [see for example [10], and [11] for a holographic approach]. FAIR and NICA are designed to explore the region of the phase diagram around and beyond the critical point, and perhaps ultimately towards the triple point. Naturally it is of great interest to have theoretical predictions as to what will happen as the chemical potential is increased in this manner. We now turn to a particular phenomenon which might be affected.

---

[2] For other possible versions, see [7].

When heavy ions collide peripherally, a very simple and compelling kinematic argument shows that, for favourable impact parameters, large quantities of angular momentum *must* be transferred to the QGP, should it be formed [12][13][14]. This is based on the simple fact that the nucleon density in each ion is non-uniform in the transverse directions, resulting in a *shearing* motion within the QGP formed in a peripheral collision. [See [13] for a particularly clear discussion.] This can be expected to have important indirect consequences [for studies of the vorticity of the plasma [15][16][17]], but it might also have directly observable implications. It has been very plausibly argued [12][13][14] that spin-orbital coupling will under these circumstances entail global quark polarization effects, leading to several striking and characteristic observable phenomena.

Thus far, no unambiguous signals of this kind have been reported by the RHIC experiments: see for example [18][19] for searches for such effects, specifically for spin alignment of certain kinds of vector mesons. Unfortunately it is very difficult [20] to estimate the precise value of the angular momentum at which observations might be expected, and there may be additional effects suppressing the polarization [21]. Becattini et al. [13] estimate that the QGP formed in collisions of gold nuclei at the RHIC might, for an optimal impact parameter, acquire an angular momentum of 7.2×10^4 in natural units, and the corresponding figure at the LHC could be significantly larger. But perhaps these values are simply not yet high enough, or collisions with optimal parameters are not yet sufficiently numerous, to produce clear-cut polarization effects. In this work we shall assume that this is the case: that, as collisions with very high angular momenta are produced in greater numbers and in different experiments, these effects will eventually be observed.

It is natural to ask whether the large chemical potentials eventually to be produced in collisions at FAIR and NICA will have any effect here. In this work, we suggest that the holographic approach [22][23][24] to the study of the QGP may give rise to an unexpected answer: *large chemical potentials tend to trigger an instability in the QGP at high angular momenta.* Presumably this would suppress polarization effects. The argument runs as follows.

As always, one begins by choosing a thermal AdS black hole spacetime, which is to be dual to a field theory representing the plasma. Just as the temperature of the QGP is dual to the Hawking temperature of the black hole, so also we expect that it should be possible to model the internal angular momentum of the QGP, as produced in peripheral collisions, by a bulk black hole endowed with angular momentum. One first thinks of the AdS-Kerr metric [25][26], and this metric has indeed been used to good effect by Atmaja and Schalm [27] to study certain aspects of a QGP endowed with large quantities of angular momentum. However, the standard AdS-Kerr metric induces a non-trivial topology [$I\!R \times \mathrm{S}^2$] on the boundary, and this boundary does not rotate differentially, in the manner decribed above. In order to obtain a more realistic geometry for the dual theory one needs to use the "rotating" [actually, *shearing*] planar black holes discovered by Klemm, Moretti, and Vanzo [28] [the "KMV_0 black holes[3]"].

---

[3] The subscript indicates planar topology for the event horizon, and also planar spatial geometry at

This apparently straightforward step, from spherical to planar topology, has an unexpected consequence, however: unlike topologically spherical AdS-Kerr black holes [29], these objects are unstable [30] when branes propagating in the bulk [as studied by Seiberg and Witten in [31]] are taken into account. This is actually a *universal* property of shearing asymptotically AdS black holes inducing planar topology and a reasonable velocity profile at infinity [32]. The brane action is in this case unbounded below, and the consequent uncontrolled brane pair-production would ultimately back-react on the black hole geometry, triggering a phase change which presumably has a dual interpretation in terms of a phase change in the plasma: the plasma ceases to exist as such, for reasons which are obscure on that side of the duality.

A key observation in [30] was that the instability studied there is only physically relevant if it can evolve during the lifetime of the QGP formed in a heavy ion collision. This lifetime is of course very short: it is estimated that local thermal equilibrium is established around 1 fm/c, and that the QGP hadronizes at around 5 fm/c after the collision [33] at the RHIC, though somewhat later at the LHC ALICE experiment [34]. In [30] we proposed a way, using holography, of identifying trends in the rate at which the instability develops. We found that the rate increases very slowly with the angular momentum density. Therefore it is plausible that the instability will not be seen under normal circumstances, that is, at relatively low angular momenta: it will not have time to develop. In fact, we shall see that this remains true even for values of the angular momentum required to observe quark polarization in current experiments such as ALICE. [At *extremely* high angular momenta, the time scale of the instability may be comparable to the normal lifetime of the plasma, but it is not clear that such values can be attained.]

This discussion was based, however, on the computations performed in [30], where it is assumed that the quark chemical potential[4] is negligible. As we noted above, that is a good approximation for most current experiments, but it will not be appropriate for the beam energy scan experiments currently under way at RHIC or to be performed at FAIR and NICA. We therefore need to ask: what is the effect of non-negligible chemical potentials on these computations? This is the subject of the present work.

The introduction of a chemical potential is modelled holographically by placing an electric charge on the bulk black hole [36][37], in addition to angular momentum. Using this new background, we can again estimate the rate at which the brane instability develops in the bulk, as a function of the black hole parameters, including the electric charge. One finds that a non-zero charge modifies the rate at which the Seiberg-Witten instability develops in these systems to a significant degree. [Note that charging an AdS black hole does not in itself necessarily or indeed usually generate this instability [38][39].]

As before, we can interpret the results holographically in terms of the rate at which an instability develops in the dual field theory. One can think of the results in the following way. Fix the angular momentum/energy density ratio at some [high] value. Then, in effect, the rate at which the "dual Seiberg-Witten instability" develops turns out to vary quite rapidly with position on the quark matter phase diagram — or, rather, on that part of it which describes the QGP.

We have computed some of these characteristic times for some interesting parameter

---

infinity.

[4] For a study of the role of the chemical potential in connection with quark polarization, see [35].

values. In accordance with [40][41][42], we do not claim that our numerical results are anything more than suggestive. What they suggest, however, is that the QGP is particularly susceptible to angular-momentum-induced instability at relatively low temperatures and moderate non-zero values of the chemical potential. The expectation is then that quark polarization effects will be suppressed in the corresponding sector of the quark matter phase diagram, even at values of the angular momentum which might otherwise be expected to render them observable. This would be of interest in itself; in a sense, it would also be an experimental probe of brane physics, since the Seiberg-Witten effect only arises because of the presence of branes in the bulk.

We begin with a discussion of the relevant black hole spacetime.

2. The QKMV$_0$ Spacetime

As is well known, in the asymptotically AdS case it is possible to find black hole spacetimes with event horizons having non-spherical topology [43]. This is fortunate, since it opens the way to considering boundary field theories defined on topologically trivial spacetimes, such as those in which actual heavy ion collisions occur. Evidently we need black holes with topologically *planar* event horizons; later, however, we will also find it necessary to make use of some results from the topologically spherical case, so we begin with non-rotating charged [AdS-Reissner-Nordström] black holes of arbitrary event horizon topology.

If the event horizon is an n-dimensional [$n \geq 2$] compact space of constant curvature k $= \{-1, 0, +1\}$ [so that, in the $k = 0$ case, we are dealing with a torus or related manifold], then the $(n + 2)$-dimensional metric with asymptotic AdS curvature $-1/L^2$ is given by

$$\begin{aligned} g(\mathrm{AdSRN}^k_{n+2}) \;=\; & -\left[\frac{r^2}{L^2} + k - \frac{16\pi M}{nV[X^k_n]r^{n-1}} + \frac{8\pi Q^2}{n(n-1)\left(V[X^k_n]\right)^2 r^{2n-2}}\right]dt^2 \\ & + \frac{dr^2}{\frac{r^2}{L^2} + k - \frac{16\pi M}{nV[X^k_n]r^{n-1}} + \frac{8\pi Q^2}{n(n-1)\left(V[X^k_n]\right)^2 r^{2n-2}}} + r^2\, d\Omega^2[X^k_n]. \end{aligned} \tag{1}$$

Here $d\Omega^2[X^k_n]$ is a metric of constant curvature $k = \{-1, 0, +1\}$ on an n-dimensional space X^k_n with [dimensionless] volume $V[X^k_n]$. With the inclusion of these volume factors [44], M and Q are the ADM mass and charge respectively. However, for calculations involving local quantities [especially the local properties of the electric field around a charged black hole], we are more interested in the densities of M and Q at the event horizon of the hole. If r_h is the value of the radial coordinate r at the event horizon, then these densities are given respectively by $M/V[X^k_n]r^n_h$ and $Q/V[X^k_n]r^n_h$.

We see that, both here and in the metric itself, M and Q only occur through the quantities $M^* = M/V[X^k_n]$, $Q^* = Q/V[X^k_n]$. Therefore r_h is itself determined by M^* and Q^*, rather than by M and Q, and so the same is true of the densities, which are given by M^*/r^n_h and Q^*/r^n_h. This means that M^* and Q^* are the basic quantities from our point of view[5]. They have the following very useful property: in the $k = 0$ case they allow us to replace the torus by a space of planar topology. We simply let the volume factor tend to infinity, along with M and Q, in such a way that the quotients M^* and Q^* remain finite. This allows us to deduce metrics for black holes with event horizons having *planar* topology. [This argument is suggested by Witten's derivation [45] of a planar black hole metric from a spherical one.]

---

[5] Note that, similarly, it is Q^* that appears in the formula for the corresponding electromagnetic potential one-form.

As we explained earlier, for our current purposes we need to extend this discussion to the case where the black hole has angular momentum as well as electric charge. From equation (1) we see that four-dimensional planar AdS-Reissner-Nordström black holes have metrics of the form

$$\mathrm{g(AdSRN_4^P)} = -\left[\frac{\mathrm{r}^2}{\mathrm{L}^2} - \frac{8\pi \mathrm{M}^*}{\mathrm{r}} + \frac{4\pi \mathrm{Q}^{*2}}{\mathrm{r}^2}\right]\mathrm{dt}^2 + \frac{\mathrm{dr}^2}{\frac{\mathrm{r}^2}{\mathrm{L}^2} - \frac{8\pi \mathrm{M}^*}{\mathrm{r}} + \frac{4\pi \mathrm{Q}^{*2}}{\mathrm{r}^2}} + \mathrm{r}^2\Big[\mathrm{d}\psi^2 + \mathrm{d}\zeta^2\Big], \quad (2)$$

where ψ and ζ are dimensionless coordinates on the plane; the mass and charge parameters M^* and Q^* are defined by a limiting process, as discussed above [so that the charge density at the horizon is $\mathrm{Q}^*/\mathrm{r}_\mathrm{h}^2$, and similarly for the energy density]. Let us now add angular momentum, described by a parameter, conventionally denoted a: it is equal to the ratio of the angular momentum of the black hole to its mass [defined again by a suitable limit]. Then we obtain the charged versions [46] of the planar black holes [or "KMV_0" black holes] discovered by Klemm et al. [28]; we call these the "QKMV_0 metrics":

$$\mathrm{g(QKMV_0)} = -\frac{\Delta_\mathrm{r}\Delta_\psi\rho^2}{\Sigma^2}\,\mathrm{dt}^2 + \frac{\rho^2}{\Delta_\mathrm{r}}\mathrm{dr}^2 + \frac{\rho^2}{\Delta_\psi}\mathrm{d}\psi^2 + \frac{\Sigma^2}{\rho^2}\Big[\omega\,\mathrm{dt} - \mathrm{d}\zeta\Big]^2, \quad (3)$$

where the asymptotic curvature is $-1/\mathrm{L}^2$, and where

$$\begin{aligned}
\rho^2 &= \mathrm{r}^2 + \mathrm{a}^2\psi^2\\
\Delta_\mathrm{r} &= \mathrm{a}^2 + \frac{\mathrm{r}^4}{\mathrm{L}^2} - 8\pi \mathrm{M}^*\mathrm{r} + 4\pi \mathrm{Q}^{*2}\\
\Delta_\psi &= 1 + \frac{\mathrm{a}^2\psi^4}{\mathrm{L}^2}\\
\Sigma^2 &= \mathrm{r}^4\Delta_\psi - \mathrm{a}^2\psi^4\Delta_\mathrm{r}\\
\omega &= \frac{\Delta_\mathrm{r}\psi^2 + \mathrm{r}^2\Delta_\psi}{\Sigma^2}\,\mathrm{a}. \quad (4)
\end{aligned}$$

Notice that (3) reduces to (2) when a = 0, just as it reduces to the KMV_0 metric when $\mathrm{Q}^* = 0$. Notice too that the metric induced on t = constant sections at the horizon has determinant equal to r_h^4; therefore, M^* and Q^* retain their interpretations as in (2), so that, in particular, the charge density at the horizon is still given by $\mathrm{Q}^*/\mathrm{r}_\mathrm{h}^2$ [though of course the numerical value has changed, because the horizon is now defined by the more complicated equation given by the vanishing of Δ_r].

With obvious changes, the properties of this geometry on and outside the horizon closely resemble those of the zero-charge KMV_0 spacetime, and the reader is referred to [28] for the details. Here we discuss only those properties directly relevant to constructing a holographic model of a dual field theory with a large angular momentum density and chemical potential.

First we note that the geometry induced on the conformal boundary is represented by a metric of the form

$$\mathrm{g(QKMV_0)_\infty} = -\mathrm{dt}^2 - \frac{2}{\gamma^2}\,\mathrm{sinl}^2\Big(\sqrt{\frac{\mathrm{a}}{\mathrm{L}}}\,\gamma\,\frac{\mathrm{x}}{\mathrm{L}}\Big)\mathrm{dtdz} + \mathrm{dx}^2 + \mathrm{dz}^2, \quad (5)$$

where $\mathrm{dx} = \mathrm{L d}\psi/\sqrt{1+\mathrm{a}^2\psi^4/\mathrm{L}^2}$, $\mathrm{dz} = \mathrm{L d}\zeta$, $\mathrm{i} = \sqrt{-1}$, $\gamma = (1+\mathrm{i})/\sqrt{2}$, and $\mathrm{sinl(x)} \equiv \mathrm{sn(x, i)}$ is the "lemniscatic sine"; it is one of the Jacobi elliptic functions [47]: see [32] for a discussion.

There are three crucial observations to be made about this boundary geometry. First, and most obviously, the geometry of the x-z plane[6] is ordinary Euclidean geometry, which of course is the actual spatial topology in which heavy-ion collisions are studied. This is obviously more realistic than the boundary topology of a topologically spherical AdS-Kerr-Newman black hole, which is also spherical in the spatial directions.

The second point to note is that while the full boundary metric is not flat, it is [globally] *conformally* flat: this was verified explicitly, though perhaps not sufficiently emphasised, in [32]. Thus, as far as conformally coupled fields are concerned, the boundary geometry is effectively Minkowskian; any misbehaviour on the part of such fields cannot be ascribed to the unusual geometry [as can happen when the boundary is not conformally flat [31]]. This is closely similar to the situation described in [48], where a rotating superconductor is described holographically using the boundary of the topologically spherical AdS-Kerr-Newman black hole: this boundary too has a non-trivial spacetime geometry, but it is [locally] conformally flat.

Thirdly, because of the off-diagonal term involving time in equation (5), zero-momentum free particles in this geometry are frame-dragged in the z direction, with a "velocity" profile which varies with x in a manner determined by the form of the lemniscatic sine. As we have seen, this sort of differential velocity profile is precisely the way in which the internal angular momentum of the actual QGP arises. Therefore, the boundary of the QKMV_0 black hole provides a model of the internal motion of the QGP in this case. [Near to the axis of the collision, the QKMV_0 profile is fairly realistic; farther away, it is not, but we would not expect the use of a fully realistic profile to change our qualitative conclusions very greatly: see [32] for the reasons.]

We stress that this result would *not* have been obtained if we had used the topologically spherical AdS-Kerr-Newman geometry. The latter does have differential rotation rates outside the event horizon, *but this effect fades away* at large distances, and the topological sphere at infinity rotates [like the event horizon] as if it were a rigid body [49].

To summarize: the transverse sections [r = constant] of the QKMV_0 black hole are not really "rotating" in the conventional sense: they are *shearing*, in the sense that the frame-dragging effect mimics a shear, and — crucially — this effect persists *even at infinity.* The QKMV_0 black hole [and related objects [32]] are therefore uniquely suited to provide a holographic description of the QGP when it is endowed with non-trivial quantities of angular momentum; which is in fact the generic case. It is clear that the QKMV_0 geometry is to be preferred, in this application, to that of the topologically spherical AdS-Kerr-Newman black hole.

Now let us turn to the question of relating the QKMV_0 parameters to those of the QGP when the latter's angular momentum density and quark chemical potential are both non-zero.

---

[6] We are following the standard conventions of the heavy-ion literature [see for example [13]]: the z axis is the axis of the collision, the x axis is transverse, and the system is analysed sectionally, that is, each y = constant plane is considered separately, so that the system is effectively two-dimensional.

3. Holography of the $\mathrm{QKMV_0}$ Parameters

The value of r at the event horizon of the $\mathrm{QKMV_0}$ black hole, $\mathrm{r_h}$, is obtained by setting Δ_r above equal to zero and taking the largest positive root:

$$\mathrm{a^2 \;+\; 4\pi Q^{*2} \;-\; 8\pi M^* r_h \;+\; r_h^4/L^2 \;=\; 0.} \tag{6}$$

We see that, as far as the position of the event horizon is concerned, the effect of adding charge is just to augment the effect of angular momentum. Quantities associated with the event horizon, such as the Hawking temperature, therefore depend on charge and angular momentum in much the same way.

As usual, the root only exists if the constant term, given by a combination of the angular momentum and charge parameters, is not too large: we have

$$\mathrm{\frac{a^2 \;+\; 4\pi Q^{*2}}{L^2} \;\leq\; 3 \times (2\pi M^*/L)^{4/3}.} \tag{7}$$

Notice that this is the only restriction on the parameter a. [This is in sharp contrast to the topologically spherical case, where, independently of the censorship condition, a is bounded by the curvature scale L, in order to ensure that frame-dragging effects do not diverge at infinity [32].]

The temperature of the black hole is given by a complicated combination of $\mathrm{M^*}$, $\mathrm{Q^*}$, and a, but for our purposes it will be more useful to use equation (6) to eliminate $\mathrm{Q^*}$ and a, so as to express the temperature in terms of $\mathrm{M^*}$ and $\mathrm{r_h}$ only [as in [28]]:

$$\mathrm{T \;=\; \frac{2r_h^3 \;-\; 4\pi M^* L^2}{2\pi L^2 r_h^2}.} \tag{8}$$

One can readily verify that this decreases monotonically as either charge or angular momentum is added to a black hole with fixed $\mathrm{M^*}$, and that it vanishes for all extremal holes. Since we will be supposing that this black hole is dual to a field theory describing a quark-gluon plasma, which of course cannot be arbitrarily cold, something must render these black holes unstable before they reach extremality; a fact also strongly suggested by the non-vanishing of the entropy in that limit, as Hartnoll pointed out in [50]. We have shown elsewhere [38][30] that the Seiberg-Witten effect can explain this, whether we approach extremality by increasing the charge or by increasing the angular momentum. Here we will study the effects of doing both, simultaneously.

Let us assume that the $\mathrm{QKMV_0}$ black hole has a dual interpretation, in the usual holographic sense. We now ask whether it is possible to reconstruct the black hole geometry, given the relevant physical parameters on the boundary.

The quantities T and a have a direct holographic interpretation in the boundary field theory: they correspond to the temperature of the field theory and to the ratio of its angular momentum density to its energy density, respectively. For any asymptotically AdS spacetime, the curvature parameter L has a physical interpretation as a *time* scale: it is the scale set by the proper time required for an object to fall from infinity to any fixed location. It should therefore be associated with some characteristic time scale of the QGP. For reasons to be explained, we assume that this scale is the "fiducial" lifetime of the QGP, that is, its lifetime in the absence of induced instabilities of the kind we shall discuss in this work.

Typical values are as follows. As mentioned, the temperature at the critical point is thought to be around 175 MeV, or rather below 1 fm^{-1} in natural units. The temperature of the QGP might take values, in future beam energy scan experiments, very roughly[7] in the range 0.5 - 2 fm^{-1}. [Bear in mind that the QGP may be able to exist at relatively low temperatures when the chemical potential is large.] In [30] we computed a rough estimate for the parameter a, in collisions with optimal impact parameter, attained in previous runs at the RHIC experiment: it is of the order 20 fm. It is not yet clear what values the beam energy scan programmes mentioned in Section 1 will produce, but, in order to be definite, we shall take it that, in the optimal collisions, $\mathrm{a} \approx 100$ fm is a reasonable order-of-magnitude estimate for future experiments. [Actually, our results are remarkably insensitive to the value of this parameter when the chemical potential is not negligible; most of the numbers we obtain in that case would change by less than 1% if $\mathrm{a} = 100$ fm were replaced by $\mathrm{a} = 20$ fm.] Finally, as we mentioned above, the QGP is thought to have a lifetime of around 5 fm/c in heavy ion collisions at currently observed [that is, low] values of the chemical potential, so this is the independent time scale here, and we adopt this as our value for L. All of these estimates are provided mainly as an aid to intuition; we make no pretence of either accuracy or precision, which would be pointless in any case under the circumstances.

If $\mathrm{Q}^* = 0$, then, if we know a, T, and L, we can completely reconstruct the geometric parameters of the dual black hole by solving the two equations (6) and (8) for the two unknowns $\mathrm{r_h}$ and M^*. But if Q^* does not vanish, then we need a third equation. This is provided by the relation between the charge parameter and the chemical potential μ. [Note that we are referring here to the quark, not the baryonic, chemical potential.] Typical values for this quantity around the critical point are estimated to be similar to the temperature; one can hope that the experiments will ultimately probe values for μ ranging between 1 and perhaps 3 or 4 fm^{-1}.

There is a standard procedure for establishing the relation between the charge parameter and μ, but here it encounters a peculiar difficulty. For a non-rotating AdS black hole, one can transform to the Euclidean domain and show that the "timelike" Killing vector vanishes at the origin of coordinates [which corresponds to the event horizon on the Lorentzian section]. This forces the time component of the gauge potential to vanish there, and that allows one to fix the additive constant which naturally occurs in that component[8]. This constant fixes the asymptotic value of the gauge potential, which in turn is identified with a fixed multiple of the chemical potential of the boundary theory. A similar argument works for the topologically spherical AdS-Kerr-Newman metric, as we will shortly see; but in that case one has to take care to apply the argument at the points where the event horizon intersects the ergosurface.

The problem here is that, for the QKMV_0 spacetime, the event horizon *never* intersects the ergosurface [28]. The proof of this statement is straightforward: with some algebraic manipulations one can write the time-time component of the metric as

---

[7] The beam scan experiments probe lower temperatures, but note that we are only interested in collisions which definitely produce a plasma, since of course only these could have a dual description in terms of a black hole.

[8] There is of course a slightly more complicated Lorentzian argument to the same effect, using the fixed points of the Killing flow.

$$\mathrm{g(QKMV_0)_{tt}} \;=\; \frac{-\Delta_\mathrm{r} \;+\; \mathrm{a}^2\Delta_\psi}{\rho^2}, \tag{9}$$

which clearly never vanishes on the event horizon if a is not zero. In order to obtain a relation between the charge parameter and the chemical potential of the field theory, we therefore need to proceed differently.

The point to note here is that μ is determined not by the full potential, but rather only by its *asymptotic* value. As one would expect from general considerations of four-dimensional black hole theory [in the absence of any kind of scalar "hair" — see [51] for that very different situation], this value can only depend on the ADM parameters of the black hole; in fact, in the topologically spherical AdS-Reissner-Nordström case, it is given simply by the black hole charge and the value of the radial coordinate at the horizon, $\mathrm{r_h}$ — which of course is determined by the ADM parameters. [Equivalently, it is given by $\mathrm{r_h}$ together with the charge density on the horizon.] It cannot, therefore, depend on the topology of the event horizon, since this is not related to the ADM parameters. Intuitively: very far from the black hole, one cannot tell, by purely local measurements, whether the r = constant, t = constant surfaces are planar or spherical. When evaluating asymptotic quantities, such as the value of a field at arbitrarily large distances from the black hole, we can therefore perform the calculation on either the topologically spherical or the planar black hole geometry, provided that we match the parameters suitably. [This would *not* be valid, of course, for global quantities such as the integral of a function over the entire spacetime, for example, an action integral, and indeed the distinction between spherical and planar event horizons is crucial for the brane action.]

Let us use this idea to find an expression for the chemical potential of a field theory dual to a shearing, electrically charged AdS black hole.

We begin, then, by discussing the duality between electric charge on the bulk black hole, and the chemical potential of the dual theory, as it arises in the context of a topologically *spherical* AdS-Kerr-Newman metric. [Thus, the setup here is precisely the charged version of the geometry used by Atmaja and Schalm in [27].] That metric is given by [25][26]

$$\mathrm{g(AdSKN)} = -\frac{\Delta_\mathrm{r}}{\rho^2}\left[\mathrm{dt} \;-\; \frac{\mathrm{a}}{\Xi}\sin^2\theta\,\mathrm{d}\phi\right]^2 \;+\; \frac{\rho^2}{\Delta_\mathrm{r}}\mathrm{dr}^2 \;+\; \frac{\rho^2}{\Delta_\theta}\mathrm{d}\theta^2 \;+\; \frac{\sin^2\theta\,\Delta_\theta}{\rho^2}\left[\mathrm{a\,dt} \;-\; \frac{\mathrm{r}^2 \;+\; \mathrm{a}^2}{\Xi}\,\mathrm{d}\phi\right]^2, \tag{10}$$

where

$$\begin{aligned}
\rho^2 &= \mathrm{r}^2 \;+\; \mathrm{a}^2\cos^2\theta \\
\Delta_\mathrm{r} &= (\mathrm{r}^2+\mathrm{a}^2)\Big(1+\frac{\mathrm{r}^2}{\mathrm{L}^2}\Big) - 2\Xi^2\mathrm{Mr} + \frac{\Xi^2\mathrm{Q}^2}{4\pi} \\
\Delta_\theta &= 1-\frac{\mathrm{a}^2}{\mathrm{L}^2}\cos^2\theta \\
\Xi &= 1-\frac{\mathrm{a}^2}{\mathrm{L}^2}.
\end{aligned} \tag{11}$$

Here M and Q are the ADM mass and charge, and a is the ratio of the ADM angular momentum to the ADM mass. [For the factors of Ξ, see [49][52].] Note that the area of the event horizon is, because of the way Ξ occurs in the metric, given by an unusual expression:

$$\mathrm{Area_h} \;=\; \frac{4\pi(\mathrm{r_h^2} \,+\, \mathrm{a^2})}{\Xi}. \tag{12}$$

The charge density at the horizon is therefore given by

$$\varsigma_{\mathrm{h}} \;=\; \frac{\Xi \mathrm{Q}}{4\pi(\mathrm{r_h^2} \,+\, \mathrm{a^2})}. \tag{13}$$

For technical reasons it will be convenient to work, until further notice, in the Euclidean regime.

The topologically spherical Euclidean AdS-Kerr-Newman gravitational instanton metric takes the form

$$\mathrm{g(EAdSKN)} = \frac{\Delta_{\mathrm{r}}^{\mathrm{E}}}{\rho_{\mathrm{E}}^2}\left[\mathrm{dt} \,+\, \frac{\mathrm{a}}{\Xi^{\mathrm{E}}}\mathrm{sin}^2\theta\,\mathrm{d}\phi\right]^2 \,+\, \frac{\rho_{\mathrm{E}}^2}{\Delta_{\mathrm{r}}^{\mathrm{E}}}\mathrm{dr}^2 \,+\, \frac{\rho_{\mathrm{E}}^2}{\Delta_{\theta}^{\mathrm{E}}}\mathrm{d}\theta^2 \,+\, \frac{\mathrm{sin}^2\theta\,\Delta_{\theta}^{\mathrm{E}}}{\rho_{\mathrm{E}}^2}\left[\mathrm{a\,dt} \,-\, \frac{\mathrm{r}^2 \,-\, \mathrm{a}^2}{\Xi^{\mathrm{E}}}\,\mathrm{d}\phi\right]^2, \tag{14}$$

where

$$\begin{aligned}
\rho_{\mathrm{E}}^2 &= \mathrm{r}^2 \,-\, \mathrm{a}^2\mathrm{cos}^2\theta \\
\Delta_{\mathrm{r}}^{\mathrm{E}} &= (\mathrm{r}^2 - \mathrm{a}^2)\Big(1 + \frac{\mathrm{r}^2}{\mathrm{L}^2}\Big) - 2(\Xi^{\mathrm{E}})^2\mathrm{Mr} - \frac{(\Xi^{\mathrm{E}})^2\mathrm{Q}^2}{4\pi} \\
\Delta_{\theta}^{\mathrm{E}} &= 1 + \frac{\mathrm{a}^2}{\mathrm{L}^2}\,\mathrm{cos}^2\theta \\
\Xi^{\mathrm{E}} &= 1 + \frac{\mathrm{a}^2}{\mathrm{L}^2}.
\end{aligned} \tag{15}$$

Here the superscript or subscript E denotes the Euclidean version of the respective quantity [both angular momentum and charge being complexified]. When cosmic censorship is satisfied, so that an event horizon exists in the Lorentzian case, the polynomial $\Delta_{\mathrm{r}}^{\mathrm{E}}$ has a positive root $\mathrm{r_0}$, and the range of r is constrained by $\mathrm{r} \geq \mathrm{r_0}$; note that, from the definition of $\Delta_{\mathrm{r}}^{\mathrm{E}}$, $\mathrm{r_0^2} > \mathrm{a^2}$, so ρ_{E}^2 is always positive. Notice that the instanton is well-defined for all values of a [including values not permitted to the Lorentzian version of this space]. For convenience let us [purely formally] define a Euclidean version of the charge density, from equation (13):

$$\varsigma_{\mathrm{h}}^{\mathrm{E}} \;=\; \frac{\mathrm{i}\,\Xi^{\mathrm{E}}\mathrm{Q}}{4\pi(\mathrm{r_0^2} \,-\, \mathrm{a^2})}; \tag{16}$$

for reasons just explained, this is well-defined for all values of a.

For a $\mathrm{QKMV_0}$ metric with given values of $\mathrm{M^*}$, $\mathrm{Q^*}$, and a, we can construct the corresponding Euclidean instanton by complexifying the time coordinate, the charge, and the angular momentum. We match this to a topologically spherical instanton as follows: take it to have the same value of a, the same value of the Euclidean radial coordinate at the origin, $\mathrm{r_0}$, and the same value of the Euclidean charge density, so that

$$\frac{\mathrm{Q^*}}{\mathrm{r_0^2}} \;=\; \frac{\Xi^{\mathrm{E}}\mathrm{Q}}{4\pi(\mathrm{r_0^2} \,-\, \mathrm{a^2})}. \tag{17}$$

This fixes the topologically spherical instanton, because equation (17) allows us to compute Q in terms of $\mathrm{M^*}$, $\mathrm{Q^*}$, and a, and then the two equations for $\mathrm{r_0}$ allow us to determine M in the same way.

Returning to the topologically spherical instanton: the Euclidean field potential is given, up to a gauge transformation, by

$$\mathrm{A^E} \;=\; \left[-\,\frac{\Xi^{\mathrm{E}}\mathrm{Qr}}{4\pi\rho_{\mathrm{E}}^2} \;+\; \kappa\right]\mathrm{dt} \;-\; \frac{\mathrm{aQr}\sin^2\!\theta}{4\pi\rho_{\mathrm{E}}^2}\mathrm{d}\phi; \tag{18}$$

here κ is a constant which could be removed by a choice of gauge, but which we shall need below. Notice that, through ρ_{E}, the coefficients here depend on both r and θ.

The Euclidean squared length of the Killing vector ∂_{t} is given by

$$|\partial_{\mathrm{t}}|^2_{\mathrm{E}} \;=\; \frac{\Delta_{\mathrm{r}}^{\mathrm{E}} \;+\; \mathrm{a}^2\sin^2\!\theta\,\Delta_{\theta}^{\mathrm{E}}}{\rho_{\mathrm{E}}^2}. \tag{19}$$

The geometry being Euclidean, this means that ∂_{t} itself must vanish when $\Delta_{\mathrm{r}}^{\mathrm{E}}$ vanishes and θ is either zero or π, and hence $\mathrm{A^E}$ must satisfy $\mathrm{A_t^E} = \mathrm{A^E}(\partial_{\mathrm{t}}) = 0$ at those locations [which correspond to the points where the ergosurface touches the event horizon in the Lorentzian version of the geometry].

Substituting $\mathrm{r} = \mathrm{r_0}$ and $\theta = 0$ or π into the potential, it follows that κ is given by $\kappa \;=\; \Xi^{\mathrm{E}}\mathrm{Qr_0}/4\pi(\mathrm{r_0^2} - \mathrm{a}^2)$, so that we have

$$\mathrm{A^E} \;=\; \left[-\,\frac{\Xi^{\mathrm{E}}\mathrm{Qr}}{4\pi\rho_{\mathrm{E}}^2} \;+\; \frac{\Xi^{\mathrm{E}}\mathrm{Qr_0}}{4\pi(\mathrm{r_0^2} - \mathrm{a}^2)}\right]\mathrm{dt} \;-\; \frac{\mathrm{aQr}\sin^2\!\theta}{4\pi\rho_{\mathrm{E}}^2}\mathrm{d}\phi. \tag{20}$$

Using equation (17) we may write this as

$$\mathrm{A^E} \;=\; \left[-\,\frac{\Xi^{\mathrm{E}}\mathrm{Qr}}{4\pi\rho_{\mathrm{E}}^2} \;+\; \frac{\mathrm{Q}^*}{\mathrm{r_0}}\right]\mathrm{dt} \;-\; \frac{\mathrm{aQr}\sin^2\!\theta}{4\pi\rho_{\mathrm{E}}^2}\mathrm{d}\phi. \tag{21}$$

If we assume that the Lorentzian version of this geometry is dual to a field theory with a non-zero chemical potential μ, then [37] μ is obtained as a multiple[9] of the asymptotic value of the time component of A. We therefore have a deceptively simple-looking result[10]:

$$\mu \;=\; \frac{\mathrm{Q}^*}{\pi\mathrm{Lr_h}}. \tag{22}$$

In accordance with the above discussion, we take it that this expression, derived in the topologically spherical setting, is also valid for the $\mathrm{QKMV_0}$ geometry.

Combining this result with equations (6) and (8) we obtain finally the following three equations:

$$4\pi\mathrm{T} \;=\; \frac{3\mathrm{r_h}}{\mathrm{L}^2} \;-\; \frac{\mathrm{a}^2}{\mathrm{r_h^3}} \;-\; \frac{4\pi^3\mu^2\mathrm{L}^2}{\mathrm{r_h}}; \tag{23}$$

$$8\pi\mathrm{M}^* \;=\; \frac{\mathrm{r_h^3}}{\mathrm{L}^2} \;+\; \frac{\mathrm{a}^2}{\mathrm{r_h}} \;+\; 4\pi^3\mathrm{L}^2\mu^2\mathrm{r_h}; \tag{24}$$

$$4\pi\mathrm{Q}^{*2} \;=\; 4\pi^3\mu^2\mathrm{r_h^2}\mathrm{L}^2. \tag{25}$$

---

[9] See [53]; we follow the same conventions.

[10] Note that the chemical potential in the topologically spherical case is given, according to this derivation, by $\mu = \Xi\mathrm{Qr_h}/[4\pi^2\mathrm{L}(\mathrm{r_h^2} + \mathrm{a}^2)]$, which is not at all formally similar to the result in the non-rotating case. In both formulae, in any case, μ depends indirectly on *all* of the black hole parameters, through $\mathrm{r_h}$.

It is elementary to show that, given positive values for T, a, and L, equation (23) always has exactly one positive real solution for $\mathrm{r_h}$. Once we have $\mathrm{r_h}$, then the remaining equations allow us to compute M^* and Q^* if we are given a, μ, and L. In other words, given the physical parameters T, a, L, and μ in the boundary field theory, we can always solve these equations for the remaining geometric parameters M^* and Q^*, and so we can find the dual $\mathrm{QKMV_0}$ bulk geometry.

We now show how to use this knowledge to investigate the evolution of the Seiberg-Witten effect in the bulk for given boundary parameters.

4. Instability of the $\mathrm{QKMV_0}$ Black Hole

The basic objects in applications of the AdS/CFT duality are the AdS-Schwarzschild black holes with either spherical or planar event horizons. These geometries can, however, be modified in a variety of interesting ways; for example, one can directly distort the geometry, as in [54][55][56], or one can distort it indirectly by adding charge or angular momentum.

Distortions of the bulk can cause the bulk theory [and therefore, by the AdS/CFT correspondence, the boundary field theory] to become unstable. Specifically, by considering a particular kind of probe brane, Seiberg and Witten [31] identified an extremely general form of instability that can arise under such circumstances, and in fact this "Seiberg-Witten instability" plays an essential role in maintaining the validity of the AdS/CFT correspondence [57].

In some cases, *but not in all*, one can see how this works explicitly, because the distortion of the bulk is sometimes accompanied by a corresponding distortion of the conformal geometry at infinity, and this affects conformal couplings to that geometry: see the discussion in [57].

This straightforward explicit description is not always possible, however, because it is certainly *not* true that every modification of the bulk is reflected in the boundary conformal structure: for example, increasing the mass of an AdS Schwarzschild black hole has no effect whatever on its boundary conformal structure, which remains locally conformally flat[11]. The same is true here: as was mentioned in Section 2 above, the boundary remains globally conformally flat when angular momentum [and charge] are added to the planar AdS black hole, and so [since the conformal structure is represented by a metric of zero scalar curvature[12]] there is *no* effect on conformally coupled fields. We shall see that the addition of angular momentum does lead to Seiberg-Witten instability in the bulk of a planar AdS black hole. By the AdS/CFT correspondence, this implies that the boundary theory must also be unstable, but this instability is clearly *not* due to any pathologies in the boundary conformal structure. This means that it is possible for the AdS/CFT correspondence to reveal an instability which might be very difficult to detect in the boundary theory itself.

Before proceeding to the computation of the relevant brane action, we have the following observation. When AdS black holes are unstable in the Seiberg-Witten sense, one finds that the effect arises initially in regions where the radial coordinate is *very large.* In more detail: the instability arises because the brane action becomes negative [and, in all of the cases to be considered here, unbounded below] beyond some value of the radial coordinate. But, if for example the angular momentum is slowly increased from zero, this happens "near infinity", *not* near the event horizon.

---

[11] See [55][56] for the mathematics of this observation.

[12] In the topologically spherical case considered by Sonner [48], the boundary is locally but not globally conformally flat, so the conformal structure is represented by a metric of positive scalar curvature.

For example, for a shearing charged planar black hole with dual parameters given by T = 2 fm^{-1}, μ = 1 fm^{-1}, a = 100 fm, and L = 5 fm, the relevant brane action [see below for the details[13]] has a graph of the kind shown in Figure 2. The action is clearly negative beyond a certain large value of r, but not near to the black hole. As the angular momentum is reduced, the curve bends down less, and the region where the action is negative moves *outward.* If the angular momentum and the charge are precisely zero, then the curve is actually asymptotic to a positive constant, so the action is positive and increasing everywhere: a brane nucleating in that geometry will tend to contract back into the black hole. But as long as the angular momentum is not zero, the system will always be unstable, in the sense that a brane which nucleates in a region *sufficiently far*



Figure 2: [Multiple of the] Brane Action,T = 2 fm^{-1}, μ = 1 fm^{-1}, a = 100 fm, L = 5 fm.

---

[13] The function being graphed here is actually the expression in braces on the right side of equation (26) below.

away will always be able to decrease its action globally [below the value near the horizon] by expanding; it can escape to infinity.

This aspect of the instability is useful, because it shows that the effect is extremely robust. For example, we cannot hope to show that couplings other than that of the brane to the metric [such as might arise, in the charged case, if one were able to obtain the electric field by means of a Kaluza-Klein reduction of some higher-dimensional theory] will remove the instability. To do so, they would have to be effective in the asymptotic region, where there should be no "hair" other than the usual ADM parameters of the black hole metric; this is highly implausible[14]. We conclude that, certainly at angular momenta and charges which are not extremely large, we can analyse the instability just by studying the effect of the spacetime geometry on the structure of the pure brane action. [This was the procedure followed, for example, by Maldacena and Maoz [58], who were the first to consider Seiberg-Witten instability in the presence of a gauge field.]

One might be concerned that this could break down at very high angular momenta or charges, but, in practice, our numerical data suggest that the instability arises at large values of the radial coordinate for all physically relevant values of these parameters; in fact, the numerical data show that, particularly in the regime of relatively low temperatures in which we are interested here, the region of negative action moves *outward* after the charge on the hole exceeds a certain [not very large] critical value. In general, then, this region is well within the zone where the only relevant parameters are the ADM parameters of the black hole. We shall therefore proceed as suggested, though a more elaborate computation may well ultimately be worthwhile.

Let us turn, then, to the form taken by the Seiberg-Witten brane action when the background is given by the $\mathrm{QKMV_0}$ geometry. Formally, the computations are precisely as in [30], though with one improvement, as follows. In that work, in order to avoid an apparent irregularity in the corresponding instanton geometry, we enforced the condition $-\Psi \leq \psi \leq \Psi$, where ψ is the Euclidean version of the coordinate which appears in equations (2) and (3), and where Ψ has to satisfy $\Psi < \sqrt{\mathrm{L/a}}$. In [32], however, it was shown that [when a is strictly non-zero, as we assume henceforth] it is possible to replace ψ with a coordinate ξ parametrising a circle, defined by

$$\psi \;=\; \sqrt{\frac{\mathrm{L}}{\mathrm{a}}}\,\mathrm{sinl}\Big(\sqrt{\frac{\mathrm{a}}{\mathrm{L}}}\,\xi\Big); \tag{26}$$

here sinl(x) is the lemniscatic sine, a periodic function ranging between ±1. Thus in fact the apparent irregularity is a coordinate problem: ψ ranges between $\pm\sqrt{\mathrm{L/a}}$, the two values corresponding to the same point. With this modification, we can simply follow the discussion in [30]. Essentially this involves a somewhat intricate computation of the volume and area of a brane inhabiting the Euclidean version of the $\mathrm{QKMV_0}$ geometry, followed by a continuation back to Lorentzian signature. Eventually we find that the brane action in this case is, for a brane of tension Θ,

$$\$[\mathrm{QKMV_0}] \;=\; \frac{2\mathrm{a}\Theta}{\mathrm{T^2 r_h^2}}\left\{\frac{\mathrm{r^2}}{2}\sqrt{\mathrm{a^2} + \frac{\mathrm{r^4}}{\mathrm{L^2}} - 8\pi \mathrm{M^* r} + 4\pi \mathrm{Q^{*2}}}\left[\frac{1}{\mathrm{a}}\mathrm{arcsinh}\Big(\frac{\sqrt{\mathrm{aL}}}{\mathrm{r}}\Big) + \frac{\sqrt{\mathrm{L/a}}}{\mathrm{r}}\sqrt{1+\frac{\mathrm{aL}}{\mathrm{r^2}}}\right]\right.$$

---

[14] Furthermore, they would have to remove an infinite quantity of negative action.

$$-\frac{1}{\mathrm{L}}\left[\sqrt{\frac{\mathrm{L}}{\mathrm{a}}}(\mathrm{r}^3-\mathrm{r}_\mathrm{h}^3)+\sqrt{\mathrm{aL}^3}(\mathrm{r}-\mathrm{r}_\mathrm{h})\right]\Bigg\}. \tag{27}$$

Note that the action vanishes at the event horizon, and that the expression is only valid at and beyond $\mathrm{r} = \mathrm{r}_\mathrm{h}$.

As we have seen, it is possible to compute r_h, M^*, and Q^* if we are given the physical data from the boundary theory [its temperature, angular momentum density, energy density, and chemical potential]. Thus, all of the parameters in the action are known in terms of those data. We can now study the behaviour of the brane action as those quantities are varied.

In all cases, however, we can write the action in the following form:

$$\begin{aligned}\$[\mathrm{QKMV}_0] \;=\; \frac{2\mathrm{a}\Theta}{\mathrm{T}^2\mathrm{r}_\mathrm{h}^2}\Bigg\{&\frac{-5\sqrt{\mathrm{aL}}}{6}\,\mathrm{r} + \left[\frac{\mathrm{r}_\mathrm{h}^3}{\sqrt{\mathrm{aL}}} + \sqrt{\mathrm{aL}}\mathrm{r}_\mathrm{h} - 4\pi\mathrm{M}^*\sqrt{\mathrm{L}^3/\mathrm{a}}\right]\\ &+ \frac{(\mathrm{a}^2+4\pi\mathrm{Q}^{*2})\sqrt{\mathrm{L}^3/\mathrm{a}}}{2\mathrm{r}} + \mathrm{O}(1/\mathrm{r}^2)\Bigg\}. \end{aligned}\tag{28}$$

For small values of r, the action is positive; but one sees from this equation that it rather quickly comes to resemble a linear function with a negative slope proportional to the square root of the parameter a [see Figure 2], which implies that, for all non-zero angular momenta, the action will become and remain *negative* beyond some value of r; it is in fact unbounded below. All of these black holes are unstable. Holographic duality then implies that *a non-zero angular momentum density is an intrinsically destabilizing influence in the dual field theory.* The effect becomes more marked at larger values of a, and so one speaks of a "holographic angular momentum cutoff". We see now that the effect is present whether the chemical potential vanishes or not, since the presence of Q^* does not change the asymptotic form of the action function [though of course it does change its numerical value at any given value of the radial coordinate]. The question is how to assess the effect of charge in a quantitative manner.

In a system evolving as rapidly as the quark-gluon plasma, instability is not necessarily a problem: all instabilities take time to evolve, and this time may not be available — a point which often plays a crucial role in heavy ion physics [see [59] for a recent example]. In [30] we proposed a simple way of measuring the time required for the Seiberg-Witten instability to develop for the KMV_0 geometry. Let us see how this extends to the QKMV_0 case.

5. Time Evolution of SW Instability in the Quark Matter Phase Plane

It is clear that the time required for the Seiberg-Witten instability to manifest itself in these systems should be related to the time required for the region of negative action to influence the entire bulk, including the region near to the event horizon.

Now AdS itself has a natural time scale, set in the following way. The proper time required for a massive point particle to fall from rest [at any point, but we can take the point to be at infinity] to the centre of AdS is given [60] by $\pi\mathrm{L}/2$. Thus L, which we normally regard as a "curvature radius", actually has a more directly physical role: it sets the time scale for dynamics in AdS and for the dual field theory.

It is natural to adapt this procedure to the present case: the time required for the entire black hole spacetime to be affected by the misbehaviour of branes lying beyond $\mathrm{r} = \mathrm{r_{neg}}$ [the value of r where the graph of the brane action crosses the horizontal axis] should be determined by the proper time required for a massive but electrically neutral particle, released from rest relative to a zero angular momentum observer, to fall from a point with $\mathrm{r} = \mathrm{r_{neg}}$ to $\mathrm{r} = \mathrm{r_h}$, the location of the event horizon.

For simplicity we consider the case of orbits in the $\mathrm{QKMV_0}$ geometry satisfying $\psi = 0$, the "equatorial" plane. Then the equations for the orbit are, as in [30],

$$\dot{\mathrm{t}}\left[\frac{-\,\Delta_{\mathrm{r}}}{\mathrm{r}^2} \;+\; \frac{\mathrm{a}^2}{\mathrm{r}^2}\right] \;-\; \dot{\zeta}\mathrm{a} \;=\; \text{constant}, \tag{29}$$

$$-\,\dot{\mathrm{t}}\mathrm{a} \;+\; \dot{\zeta}\mathrm{r}^2 \;=\; 0, \tag{30}$$

where the dot denotes a derivative with respect to proper time. Combining these with the fact that the geodesic tangent is of unit length and with the initial conditions, one finds that the time required, which we denote by $\tau(\mathrm{a},\, \mathrm{T},\, \mu,\, \mathrm{L})$, is given by

$$\tau(\mathrm{a},\mathrm{T},\mu,\mathrm{L}) \;=\; \int_{\mathrm{r_h}}^{\mathrm{r_{neg}}}\left[\frac{\mathrm{a}^2+4\pi\mathrm{Q}^{*2}}{\mathrm{r_{neg}^2}} + \frac{\mathrm{r_{neg}^2}}{\mathrm{L}^2} - \frac{8\pi\mathrm{M}^*}{\mathrm{r_{neg}}} - \frac{\mathrm{a}^2+4\pi\mathrm{Q}^{*2}}{\mathrm{r}^2} - \frac{\mathrm{r}^2}{\mathrm{L}^2} + \frac{8\pi\mathrm{M}^*}{\mathrm{r}}\right]^{-1/2}\mathrm{dr}. \tag{31}$$

Given the field theory parameters a, T, L, and μ, we can compute, as we have discussed, $\mathrm{r_h}$, M^*, and Q^*, so $\mathrm{r_{neg}}$ can be computed by finding the larger positive real root of the expression in equation (27). Thus everything in this expression is now known to us in terms of the boundary data.

Let us examine some explicit values, treating them with due caution. In accordance with our earlier discussion, we take a to be around 100 fm [though, as we mentioned earlier, the results vary with a only slightly when the chemical potential is non-zero]. To fix L, recall from our discussion above that when the angular momentum/energy density ratio and the chemical potential are small, as they have been in experiments performed hitherto, then the point where the action graph cuts the horizontal axis recedes to infinity. Letting $\mathrm{r_{neg}}$ tend to infinity[15] in equation (31), we find that the time required is again $\pi\mathrm{L}/2$, so, as above, we take it that L is the characteristic time scale defined by the plasma as it appears in these experiments, roughly 5 fm/c.

We can now numerically investigate $\tau(\mathrm{a},\, \mathrm{T},\, \mu,\, \mathrm{L})$ as a function of position in the quark matter phase diagram, specifically in the region $0.5 < \mathrm{T} < 2$, $0 < \mu < 5$ in units of inverse fm. The results are shown in the table: the times are given as multiples of the approximate lifetime of the QGP in heavy ion collisions, so that values below unity mean that the instability can develop within the normal lifetime of the plasma[16].

	$\mu = 0$	$\mu = 1$	$\mu = 2$	$\mu = 3$	$\mu = 4$	$\mu = 5$
$\mathrm{T} = 2$	1.5522	1.5169	1.1125	0.91000	0.81236	0.75637
$\mathrm{T} = 1$	1.5006	1.0975	0.81091	0.71996	0.67685	0.65187
$\mathrm{T} = 1/2$	1.3370	0.80525	0.67613	0.63535	0.61558	0.60310

---

[15] This is most easily done by changing to a new variable, $\mathrm{s} = \mathrm{r}/\mathrm{r_{neg}}$, to replace r, though care in taking the limit should be exercised.

[16] We remind the reader that the $\mathrm{T} = 1/2$, $\mu = 0$ entry is not physical, because the plasma does not exist in that corner of the phase diagram. The same may be true of the $\mathrm{T} = 1/2$, $\mu = 1$ entry.

Taken at face value, the numbers indicate that, when the chemical potential is small, the time required for the Seiberg-Witten instability to develop is considerably longer than the time available: the QGP can evolve without interference in that case. But as one scans across the table to larger values of the chemical potential, the instability develops at a dramatically enhanced rate[17], effectively shortening the lifetime of the plasma. Note that quark polarization is thought [14] to be a cumulative process, building up as a result of multiple scattering; therefore it requires a substantial period of time, relative to the lifetime of the plasma, to reach observable levels, if it does so. A reduction of the plasma lifetime by the large factors indicated here can therefore be expected to have important consequences.

6. Conclusion: Quark Polarization in the Phase Diagram

There are at least two ways of using the numbers in the table above. First, if one begins in the region of the quark matter phase diagram currently being investigated by the ALICE experiment [34] at the LHC [large T, $\mu \approx 0$] and moves diagonally down to around T = 1 fm^{-1}, $\mu = 3\ \text{fm}^{-1}$, one sees that the time required for the instability to develop falls quite dramatically; so if we observe quark polarization effects in the first case, and then scan to large values of the chemical potential, we might find that the rate at which such events are seen would decrease accordingly. Secondly, we see that this reduction might be even more marked if one scanned down to still lower temperatures but more moderate values of μ, say to T = 1/2 fm^{-1}, $\mu = 2\ \text{fm}^{-1}$ — that is, to a region of the phase diagram which might realistically be accessible to FAIR and NICA. If one could see both, then an explanation in terms of the Seiberg-Witten effect might begin to seem plausible. [In fact, the numbers suggest that it is precisely for μ in the range of 1 to 2 or 3 fm^{-1} that the reduction in the times is most marked; there is little to be gained by going to 4 or 5 fm^{-1}, even if that were possible.]

There are at least two problems with this scenario, one theoretical, the other observational. First, we have been assuming that the parameter a remains fixed. Since a is the *ratio* of the angular momentum density to the energy density, it is not clear exactly how it varies during scans of the sort we are describing, since both densities will decrease, on average, for collisions at lower energies; one expects it to fall somewhat, though not very drastically. As we have mentioned, in practice this has very little effect on the numbers we obtain, so this is not a matter for concern. The second difficulty is more serious: precisely because the scan reduces the energies of the collisions, high angular momentum events will be harder to identify in any case, and one would need to separate this from the effect we have been describing. In addition, there is some reason for concern that the hydrodynamic approximation may start to break down in collisions of this kind: see the data reported in [62].

Even as a qualitative prediction, then, the idea that a "holographic" abbreviation of the plasma lifetime may be observed as experiments scan to large values of the quark chemical potential seems optimistic, for reasons we have just discussed, in addition to all of the usual reasons: QCD is not strongly coupled in the UV, its coupling runs, and so on. Nevertheless it is useful to ask what it would mean if something of this sort were observed. The AdS/CFT correspondence is often described as a "gauge-gravity" duality, but [40] this is a misnomer: it is important for the duality that there are other objects in the bulk, apart from the metric. Seiberg-Witten instability reveals a concrete aspect of this, since it involves branes in a basic way. Any experimental evidence, however imprecise, suggesting that this effect is real, would therefore also be direct evidence that the correct dual description of QCD involves branes, in an equally basic way.

---

[17] In fact it is possible that we are overestimating the times here, since lower temperatures in themselves tend to increase rates of brane nucleation: see [61] [though note that the calculations done there assume an event horizon with the topology of a compact hyperbolic space].

Acknowledgement

The author is very grateful for the kind hospitality of NORDITA, for providing facilities and an environment in which this work could be done, and wishes to thank particularly Lárus Thorlacius and the organisers of the "Holographic Way" programme. He also wishes to thank Soon Wanmei for support, encouragement, and advice on numerical questions.

References

[1] Sourendu Gupta, Xiaofeng Luo, Bedangadas Mohanty, Hans Georg Ritter, Nu Xu, Scale for the Phase Diagram of Quantum Chromodynamics, Science 332 (2011) 1525, arXiv:1105.3934 [hep-ph]

[2] Akira Ohnishi, Phase diagram and heavy-ion collisions: Overview, Prog.Theor.Phys.Suppl. 193 (2012) 1, arXiv:1112.3210 [nucl-th]

[3] Ilya Selyuzhenkov, Recent experimental results from the relativistic heavy-ion collisions at LHC and RHIC, arXiv:1109.1654 [nucl-ex]

[4] Xin Dong (for the STAR Collaboration), Highlights from STAR, arXiv:1210.6677 [nucl-ex]

[5] M. Bleicher, M. Nahrgang, J. Steinheimer, P. Bicudo, Physics Prospects at FAIR, Acta Phys.Polon. B43 (2012) 731, arXiv:1112.5286 [hep-ph]

[6] V. D. Kekelidze, A. D. Kovalenko, I. N. Meshkov, A. S. Sorin, G. V. Trubnikov, NICA at JINR: New prospects for exploration of quark-gluon matter, Physics of Atomic Nuclei 75 (2012) 542

[7] Kenji Fukushima, Chihiro Sasaki, The phase diagram of nuclear and quark matter at high baryon density, arXiv:1301.6377 [hep-ph]

[8] Zoltan Fodor, Sandor D. Katz, Christian Schmidt, The Density of states method at non-zero chemical potential, JHEP 0703:121,2007, arXiv:hep-lat/0701022

[9] G. Endrodi, Z. Fodor, S.D. Katz, K.K. Szabo, The QCD phase diagram at nonzero quark density, JHEP 1104 (2011) 001, arXiv:1102.1356 [hep-lat]

[10] A. Andronic, D. Blaschke, P. Braun-Munzinger, J. Cleymans, K. Fukushima, L.D. McLerran, H. Oeschler, R.D. Pisarski, K. Redlich, C. Sasaki, H. Satz, J. Stachel, Hadron Production in Ultra-relativistic Nuclear Collisions: Quarkyonic Matter and a Triple Point in the Phase Diagram of QCD, Nucl.Phys. A837 (2010) 65, 0911.4806 [hep-ph]

[11] Jan de Boer, Borun D. Chowdhury, Michal P. Heller, Jakub Jankowski, A holographic realization of the Quarkyonic phase, arXiv:1209.5915 [hep-th]

[12] Zuo-Tang Liang, Xin-Nian Wang, Globally polarized quark-gluon plasma in non-central A+A collisions, Phys.Rev.Lett. 94 (2005) 102301, Erratum-ibid. 96 (2006) 039901, arXiv:nucl-th/0410079

[13] F. Becattini, F. Piccinini, J. Rizzo, Angular momentum conservation in heavy ion collisions at very high energy, Phys.Rev.C77:024906,2008, arXiv:0711.1253 [nucl-th]

[14] Xu-Guang Huang, Pasi Huovinen, Xin-Nian Wang, Quark Polarization in a Viscous Quark-Gluon Plasma, Phys. Rev. C84, 054910(2011), arXiv:1108.5649 [nucl-th]

[15] Victor Roy, A. K. Chaudhuri, Bedangadas Mohanty, Comparison of results from a 2+1D relativistic viscous hydrodynamic model to elliptic and hexadecapole flow of charged hadrons measured in Au-Au collisions at $\sqrt{s_{\rm NN}} = 200$ GeV, Phys.Rev. C86 (2012) 014902, arXiv:1204.2347 [nucl-th]

[16] Marco M. Caldarelli, Robert G. Leigh, Anastasios C. Petkou, P.Marios Petropoulos, Valentina Pozzoli, Konstadinos Siampos, Vorticity in holographic fluids, PoS CORFU2011 (2011) 076, arXiv:1206.4351

[17] Mircea Baznat, Konstantin Gudima, Alexander Sorin, Oleg Teryaev, Helicity separation in Heavy-Ion Collisions, arXiv:1301.7003 [nucl-th]

[18] J H Chen (for the STAR Collaboration), Spin alignment of $K^{*0}(892)$ and ϕ (1020) mesons in Au+Au and p+p collisions at $\sqrt{s_{\rm NN}} = 200$ GeV, J. Phys. G: Nucl. Part. Phys. 35 (2008) 044068

[19] STAR Collaboration (B.I. Abelev et al.), Spin alignment measurements of the $K^{*0}(892)$ and ϕ (1020) vector mesons in heavy ion collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys.Rev. C77 (2008) 061902, arXiv:0801.1729 [nucl-ex]

[20] Jian-Hua Gao, Shou-Wan Chen, Wei-tian Deng, Zuo-Tang Liang, Qun Wang, Xin-Nian Wang, Global quark polarization in non-central A+A collisions, Phys.Rev. C77 (2008) 044902, arXiv:0710.2943 [nucl-th]

[21] Alejandro Ayala, Eleazar Cuautle, G. Herrera Corral, J. Magnin, Luis Manuel Montano, Spin alignment of vector mesons in heavy ion and proton - proton collisions, Phys.Lett. B682 (2010) 408, arXiv:0906.4295 [hep-ph]

[22] Jorge Casalderrey-Solana, Hong Liu, David Mateos, Krishna Rajagopal, Urs Achim Wiedemann, Gauge/String Duality, Hot QCD and Heavy Ion Collisions, arXiv:1101.0618 [hep-th]

[23] Mariano Chernicoff, J. Antonio Garcia, Alberto Guijosa, Juan F. Pedraza, Holographic Lessons for Quark Dynamics, J.Phys.G G39 (2012) 054002, arXiv:1111.0872 [hep-th]

[24] Youngman Kim, Ik Jae Shin, Takuya Tsukioka, Holographic QCD: Past, Present, and Future, arXiv:1205.4852 [hep-ph]

[25] B. Carter, Hamilton-Jacobi and Schrodinger separable solutions of Einstein's equations, Commun.Math.Phys. 10 (1968) 280

[26] S.W. Hawking, C.J. Hunter, Marika Taylor, Rotation and the AdS/CFT correspondence, Phys.Rev.D59:064005,1999, arXiv:hep-th/9811056

[27] A. Nata Atmaja, K. Schalm, Anisotropic Drag Force from 4D Kerr-AdS Black Holes, JHEP 1104:070,2011, arXiv:1012.3800 [hep-th]

[28] D. Klemm, V. Moretti, L. Vanzo, Rotating Topological Black Holes, Phys.Rev.D57:6127,1998; Erratum-ibid.D60:109902,1999, arXiv:gr-qc/9710123

[29] Brett McInnes, Kerr Black Holes Are Not Fragile, Nucl. Phys. B 857 (2012) 362, arXiv:1108.6234 [hep-th]

[30] Brett McInnes, Fragile Black Holes and an Angular Momentum Cutoff in Peripheral Heavy Ion Collisions, Nucl. Phys. B861 (2012) 236, arXiv:1201.6443 [hep-th]

[31] Nathan Seiberg, Edward Witten, The D1/D5 System And Singular CFT, JHEP 9904 (1999) 017, arXiv:hep-th/9903224

[32] Brett McInnes, Universality of the Holographic Angular Momentum Cutoff, Nucl.Phys. B864 (2012) 722, arXiv:1206.0120 [hep-th]

[33] Andreas Ipp, Antonino Di Piazza, Jorg Evers, Christoph H. Keitel, Photon polarization as a probe for quark-gluon plasma dynamics, Phys.Lett. B666 (2008) 315, arXiv:0710.5700 [hep-ph]

[34] ALICE Collaboration (Klaus Reygers for the collaboration), Quick Tour of Ultra-Relativistic Heavy-Ion Physics at the LHC, arXiv:1208.1626 [nucl-ex]

[35] Jian-Hua Gao, Zuo-Tang Liang, Shi Pu, Qun Wang, Xin-Nian Wang, Chiral Anomaly and Local Polarization Effect from Quantum Kinetic Approach, arXiv:1203.0725 [hep-ph]

[36] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, Robert C. Myers, Thermodynamics and fluctuations of charged AdS black holes, Phys.Rev. D60 (1999) 104026, hep-th/9904197

[37] Shinpei Kobayashi, David Mateos, Shunji Matsuura , Robert C. Myers , Rowan M. Thomson, Phase transitions at finite baryon density, JHEP 0702 (2007) 016, hep-th/0611099 [hep-th]

[38] Brett McInnes, Bounding the Temperatures of Black Holes Dual to Strongly Coupled Field Theories on Flat Spacetime, JHEP09(2009)048, arXiv:0905.1180 [hep-th]

[39] Brett McInnes, A Universal Lower Bound on the Specific Temperatures of AdS-Reissner-Nordstrom Black Holes with Flat Event Horizons, Nucl.Phys.B848 (2011) 474, arXiv:1012.4056 [hep-th]

[40] David Mateos, Gauge/string duality applied to heavy ion collisions: Limitations, insights and prospects, J.Phys.G G38 (2011) 124030, arXiv:1106.3295 [hep-th]

[41] Andreas Karch, Recent Progress in Applying Gauge/Gravity Duality to Quark-Gluon Plasma Physics, AIP Conf.Proc. 1441 (2012) 95, arXiv:1108.4014 [hep-ph]

[42] Yuri V. Kovchegov, CFT applications to relativistic heavy ion collisions: a brief review, arXiv:1112.5403 [hep-ph]

[43] J.P.S. Lemos, Phys.Lett.B353:46,1995, arXiv:gr-qc/9404041; R.B. Mann, Class.Quant.Grav. 14 (1997) L109, arXiv:gr-qc/9607071; Rong-Gen Cai, Yuan-Zhong Zhang, Phys.Rev.D54:4891,1996, arXiv:gr-qc/9609065; Danny Birmingham, Class.Quant.Grav. 16 (1999) 1197, arXiv:hep-th/9808032

[44] Dieter R. Brill, Jorma Louko, Peter Peldan, Thermodynamics of (3+1)-dimensional black holes with toroidal or higher genus horizons, Phys.Rev. D56 (1997) 3600, arXiv:gr-qc/9705012

[45] Edward Witten, Anti-de Sitter space, thermal phase transition, and confinement in gauge theories, Adv.Theor.Math.Phys.2:505,1998, arXiv:hep-th/9803131

[46] Natxo Alonso-Alberca, Patrick Meessen, Tomas Ortin, Class.Quant.Grav. 17 (2000) 2783, arXiv:hep-th/0003071; Jose P.S. Lemos, arXiv:gr-qc/0011092; A. Khodam-Mohammadi, arXiv:0707.3515 [hep-th]; Marco Astorino, JHEP 1206 (2012) 086, arXiv:1205.6998 [gr-qc]

[47] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions With Formulas, Graphs, and Mathematical Tables*, Dover Publications, New York, 1964

[48] Julian Sonner, A Rotating Holographic Superconductor, Phys.Rev.D80:084031,2009, arXiv:0903.0627 [hep-th]

[49] Marco M. Caldarelli, Guido Cognola, Dietmar Klemm, Thermodynamics of Kerr-Newman-AdS Black Holes and Conformal Field Theories, Class.Quant.Grav. 17 (2000) 399, arXiv:hep-th/9908022

[50] Sean A. Hartnoll, Lectures on holographic methods for condensed matter physics, Class.Quant.Grav.26:224002,2009, arXiv:0903.3246 [hep-th]

[51] Yen Chin Ong, Pisin Chen, Stringy Stability of Charged Dilaton Black Holes with Flat Event Horizons, JHEP 1208 (2012) 079, arXiv:1205.4398 [hep-th]

[52] G.W. Gibbons, M.J. Perry, C.N. Pope, The First Law of Thermodynamics for Kerr-Anti-de Sitter Black Holes, Class.Quant.Grav.22:1503-1526,2005, arXiv:hep-th/0408217

[53] Robert C. Myers, Miguel F. Paulos, Aninda Sinha, Holographic Hydrodynamics with a Chemical Potential, JHEP 0906:006,2009, arXiv:0903.2834

[54] Keiju Murata, Tatsuma Nishioka, Norihiro Tanahashi, Warped AdS_5 Black Holes and Dual CFTs, Prog.Theor.Phys.121:941,2009, arXiv:0901.2574 [hep-th]; Yves Brihaye, Jutta Kunz, Eugen Radu, From black strings to black holes: Nuttier and squashed AdS(5) solutions, JHEP 0908:025,2009, arXiv:0904.1566 [gr-qc]; Yves Brihaye, Eugen Radu, Cristian Stelea, Charged squashed black holes with negative cosmological constant in odd dimensions, Phys.Lett. B709 (2012) 293, arXiv:1201.3098 [hep-th]

[55] Michael T. Anderson, Piotr T. Chrusciel, Erwann Delay, Nontrivial, static, geodesically complete, vacuum space-times with a negative cosmological constant, JHEP 0210:063,2002, arXiv:gr-qc/0211006

[56] M. Anderson, P.T. Chrusciel, E. Delay, Non-trivial, static, geodesically complete space-times with a negative cosmological constant II. $n \geq 5$, Proceedings of the Strasbourg Meeting on AdS-CFT correspondence, O.Biquard, V.Turaev Eds., IRMA Lectures in Mathematics and Theoretical Physics, de Gruyter, Berlin, New York, arXiv:gr-qc/0401081

[57] Brett McInnes, Fragile Black Holes, Nucl.Phys. B842 (2011) 86, arXiv:1008.0231 [hep-th]

[58] Juan Maldacena, Liat Maoz, Wormholes in AdS, JHEP 0402 (2004) 053, arXiv:hep-th/0401024

[59] Andreas Ipp, Unstable dynamics of Yang-Mills fields at early times of heavy ion collisions, arXiv:1210.5150 [hep-ph]

[60] G.W. Gibbons, Anti-de-Sitter spacetime and its uses, Proceedings of the 2nd Samos Meeting on Cosmology, Geometry and Relativity, S Cotsakis and G W Gibbons eds, Lecture Notes in Physics 537 (2000), arXiv:1110.1206 [hep-th]

[61] Jose L.F. Barbon, Javier Martinez-Magan, Spontaneous fragmentation of topological black holes, JHEP 1008:031,2010, arXiv:1005.4439 [hep-th]

[62] Ulrich W. Heinz, Chun Shen, Huichao Song, The viscosity of quark-gluon plasma at RHIC and the LHC, AIP Conf.Proc. 1441 (2012) 766, arXiv:1108.5323 [nucl-th]